



Ultra Sensitive Oestradiol & Testosterone Anti-Müllerian Hormone (AMH)

Internal Testes or remnant testicular tissue

- Single Ultra-Sensitive Testosterone for whole testes.
- AMH Single Sensitive test to detect testicular tissue.
- hCG stimulation test (2 x testosterone) for testicular tissue.

Spayed bitch coming into season? Rescue cat - has it been spayed?

- Single Ultra-Sensitive Oestradiol (E2) if there are signs of season at the time of sampling.
- AMH Single Sensitive test to detect ovarian tissue and may be more reliable than performing the GnRH stimulation test when no signs of oestrus are present at the time of sampling.
- GnRH stimulation test (2 x E2) to detect ovarian tissue.
- hCG stimulation test (2 x E2) to detect ovarian tissue in dogs.
- hCG stimulation test (2 x Progesterone) for cats showing signs of oestrus.

Sertoli cell tumour in a male dog?

- Single sample for Ultra-Sensitive Oestradiol.

**We use highly accurate methods
to measure levels that many analysers
cannot detect!**



For further information contact
NationWide Specialist Laboratories
Unit 2, Sawston Park, London Road,
Pampisford, Cambridge, CB22 3EE
T. 01223 493400 F. 01223 493404
email: info@cslabs.co.uk or visit www.thehormonelab.com